

AperTO - Archivio Istituzionale Open Access dell'Università di Torino

Nuclear Ca²⁺ signalling in arbuscular mycorrhizal and actinorhizal endosymbioses: on the trail of novel underground signals

This is a pre print version of the following article:

Original Citation:

Availability:

This version is available <http://hdl.handle.net/2318/1633388> since 2017-05-12T17:54:54Z

Published version:

DOI:10.1111/nph.14350

Terms of use:

Open Access

Anyone can freely access the full text of works made available as "Open Access". Works made available under a Creative Commons license can be used according to the terms and conditions of said license. Use of all other works requires consent of the right holder (author or publisher) if not exempted from copyright protection by the applicable law.

(Article begins on next page)

This is the author's final version of the contribution published as:

Barker, David G; Chabaud, Mireille; Russo, Giulia; Genre, Andrea. Nuclear Ca²⁺ signalling in arbuscular mycorrhizal and actinorhizal endosymbioses: on the trail of novel underground signals. *NEW PHYTOLOGIST*. 214 (2) pp: 533-538.

DOI: 10.1111/nph.14350

The publisher's version is available at:

<http://onlinelibrary.wiley.com/doi/10.1111/nph.14350/fullpdf>

When citing, please refer to the published version.

Link to this full text:

<http://hdl.handle.net/>



New Phytologist

Nuclear Ca²⁺ signaling in arbuscular mycorrhizal and actinorhizal endosymbioses: on the trail of novel underground signals

Journal:	<i>New Phytologist</i>
Manuscript ID	NPH-TI-2016-22341
Manuscript Type:	TI – Commissioned material – Tansley insight
Date Submitted by the Author:	30-Jun-2016
Complete List of Authors:	Barker, David; UMR CNRS-INRA 2594/441, Laboratory of Plant-Microbe Interactions Chabaud, Mireille; UMR CNRS-INRA 2594/441, Laboratory of Plant-Microbe Interactions Russo, Giulia; Università di Torino, Dipartimento di Scienze della Vita e Biologia dei Sistemi Genre, Andrea; Università di Torino, Dipartimento di Scienze della Vita e Biologia dei Sistemi
Key Words:	Cameleon calcium reporters, Chitin oligomers, Common symbiotic signaling pathway, LysM receptor-like kinases, Nuclear calcium spiking, Plant-microbe interactions, Root endosymbioses

SCHOLARONE™
Manuscripts

Nuclear Ca²⁺ signaling in arbuscular mycorrhizal and actinorhizal endosymbioses: on the trail of novel underground signals

David G. Barker¹, Mireille Chabaud¹, Guilia Russo² and Andrea Genre²

¹Laboratory of Plant–Microbe Interactions (LIPM), INRA-CNRS-Toulouse University, 31326 Castanet-Tolosan, France; ²Department of Life Sciences and Systems Biology, Turin University, Viale Mattioli 25, 10125, Turin, Italy

Author for correspondence:

David G. Barker

Tel (33).5.61.28.54.15

David.Barker@toulouse.inra.fr

Word counts:

Summary: **123** words

Main Text: **2,495** words

No. Figures: **2** (Fig. 1 in colour)

No. Tables: **0**

Contents list:

Summary

I. Introduction

II. Nuclear calcium signaling and the AM symbiosis

III. Pre-infection signaling during *Frankia*/actinorhizal plant nodulation

IV. Conclusions & future outlook

Acknowledgements

References

1 **Summary**

2 Root endosymbioses are beneficial associations formed between terrestrial plants and either
 3 bacterial or fungal microorganisms. A common feature of these intracellular symbioses is the
 4 requirement for mutual recognition between the two partners prior to host-regulated microbial
 5 entry. Specific microbial factors activate a highly conserved plant signal transduction
 6 pathway, of which a central component is the triggering of sustained Ca^{2+} oscillations in the
 7 host epidermis. This then leads to the specialized cellular reprogramming required for the
 8 construction of the transcellular apoplastic microbial entry compartments. Here we focus on
 9 recent findings concerning this crucial Ca^{2+} -dependent signaling step for endosymbiotic
 10 associations involving either arbuscular mycorrhizal fungi or nitrogen-fixing *Frankia*
 11 actinomycetes, as well as how this knowledge is contributing to the identification of the
 12 respective microbial factors.

13 **Key Words:**

14 Cameleon calcium reporters; Chitin oligomers; Common symbiotic signaling pathway; LysM
 15 receptor-like kinases; Nuclear calcium spiking; Plant-microbe interactions; Root
 16 endosymbioses

I. Introduction

Throughout the evolution of land plants, mutualistic fungal and bacterial associations have provided key metabolites (phosphorus, nitrogen, microelements etc.) to their respective hosts, and thereby played a major role in plant colonization of terrestrial ecosystems. In return, the microbial partners benefited from both a source of photosynthates as well as privileged ecological niches. Striking examples of such beneficial associations are the so-called root endosymbioses, where the microsymbionts are housed within specialized host cell compartments, whether in the inner root cortex for the ancient and widespread arbuscular mycorrhizal (AM) symbiosis, or within *de novo* constructed root organs (nodules) in the case of the more recently evolved rhizobial/legume and *Frankia*/actinorhizal plant nitrogen-fixing symbioses.

Studies initially focused on the rhizobial/legume symbiosis using model legumes such as *Medicago truncatula* and *Lotus japonicus* revealed that the successful establishment of this association requires host recognition of specific rhizobial lipo-chitooligosaccharide (LCO) signals known as Nod factors (NFs; Dénarié & Cullimore, 1993). These NF LCOs are perceived *via* legume receptor-like kinases (RLK) belonging to the chitin-binding LysM-RLK family. This then activates a specific host signal transduction pathway in target root hairs, a central feature of which is the triggering of sustained nuclear-associated Ca^{2+} oscillations (known as spiking) which are decoded by a dedicated calcium and calmodulin kinase (CCaMK) (Oldroyd & Downie, 2006). Major cellular reprogramming in the host cells is thus initiated, resulting in the progressive construction of the transcellular compartment (infection thread) within root hairs, through which the rhizobia are conveyed across the root outer tissues (e.g. Fournier *et al.*, 2008). This sophisticated mode of apoplastic root penetration is thought to allow selectivity and regulation of microbial access to inner root tissues.

Legume-based research was also instrumental in revealing striking similarities between the molecular and cellular mechanisms of both rhizobial and AM fungal root colonization, including the mode of fungal root entry *via* apoplastic intracellular compartments (Genre *et al.*, 2005) as well as the activation of the Ca^{2+} spiking/CCaMK core signaling module (Kosuta *et al.*, 2008; Chabaud *et al.*, 2011) which lies at the heart of the so-called common symbiosis signaling pathway (CSSP; see Fig.1). More recently, a key role for Ca^{2+} signaling has also been demonstrated for the nitrogen-fixing endosymbioses formed between filamentous *Frankia* and their actinorhizal hosts (Chabaud *et al.*, 2015; Granqvist *et al.*, 2015). In this

review we will focus on the latest discoveries which throw light on Ca^{2+} signaling during the establishment of both the AM and *Frankia*/actinorhizal root symbioses as well as the resulting approaches which are now being employed to identify the corresponding microbial factors which activate the Ca^{2+} -dependent CSSP.

II. Nuclear calcium signaling and the AM symbiosis belonging to Glomeromycotina

AM fungi, collectively known as Glomeromycota, are widespread obligate biotrophs which are able to colonize roots of the majority (approx. 80%) of plant species, forming elaborate ramified symbiotic structures known as arbuscules within inner cortical cells. Fossil evidence has revealed that analogous structures were present in the tissues of early land plants (over 400 million years ago), suggesting that the AM symbiosis played a central role in facilitating plant access to nutrients in a harsh terrestrial environment (Bonfante & Genre, 2008). Furthermore, recent phylogenomic studies have shown that the complete CSSP module is present throughout extant plant clades, thus emphasizing the importance of microbe-host Ca^{2+} -dependent signaling even during these earliest mutualistic associations (Delaux *et al.*, 2015).

An important advance in the study of oscillatory nuclear Ca^{2+} signaling during the initial stages of endosymbiotic associations was the development of *in vivo* cameleon-based calcium reporters coupled with confocal microscopy imaging (Miwa *et al.*, 2006). In particular, the development of nuclear-localized cameleons such as Nup-YC2.1 (Sieberer *et al.*, 2009) greatly facilitated the detection of Ca^{2+} spiking in atrichoblasts, the non root hair epidermal cells which are the primary targets of AM colonization. Experiments using both legume (*M. truncatula*) and non-legume (*Daucus carota*) root organ cultures (ROCs) expressing Nup-YC2.1 were instrumental in first demonstrating nuclear Ca^{2+} spiking in atrichoblasts associated with AM fungal contact and hyphopodium formation (Chabaud *et al.*, 2011; Fig. 2a). Significantly, spiking frequency was highest in those cells where the nucleus had migrated to the site of fungal attachment, a key event which precedes the construction of transcellular apoplastic compartment (Genre *et al.*, 2005).

The presence of symbiotic fungal signals in germinating spore exudates of several AM species was also examined using Ca^{2+} spiking responses as a bio-assay in both *Medicago* and carrot ROCs expressing Nup-YC2.1. These studies led to the identification of short-chain chitin oligomers (chitotetraose and chitopentaose) as candidate AM signals (Genre *et al.*, 2013). Not only can these so-called Myc-COs activate the host CSSP at low concentrations

(10⁻⁸ M) but their levels are greatly enhanced in the spore exudates if the synthetic strigolactone GR24 is present during spore germination. Since plant strigolactones stimulate AM hyphal development prior to initial root contact, these findings provide evidence for reciprocal molecular signaling between host and fungal symbiont during the pre-infection stage. Finally, the nuclear Ca²⁺ spiking elicited in *M. truncatula* atrichoblasts by AM fungal contact, fungal exudates or short-chain Myc-COs (Fig. 2a-c) is generally less regular in both periodicity and spike profile compared to NF-activated spiking in root hairs (Russo *et al.*, 2013; Fig. 2d). The reason for this difference in spiking signature is currently unclear, and contrasts with the similar Ca²⁺ spiking profiles associated with both rhizobial and AM infection of cortical cells (Sieberer *et al.*, 2012).

In parallel to these studies, the use of NF bioassays (root hair deformation and early nodulin gene expression) had revealed other potential AM fungal signals in the form of either sulphated or non-sulphated LCOs, present in both AM spore and colonized root exudates (Maillet *et al.*, 2011). These Myc-LCOs structurally resemble rhizobial LCOs and elicit similar Ca²⁺ spiking responses to NFs in root hairs of *M. truncatula* seedlings (Sun *et al.*, 2015). Whilst *M. truncatula* mutants defective in the LysM receptor-like kinase NFP (Nod Factor Perception) fail to nodulate and are totally unresponsive to NFs (Ben Amor *et al.*, 2003), these same mutants exhibit normal AM colonization. Nevertheless, Ca²⁺ spiking is blocked in *nfp* mutants in response to exogenous Myc-LCOs (Sun *et al.*, 2015). Furthermore, transcriptomic approaches have shown that root gene expression in young seedlings in response to Myc-LCOs is also essentially dependent on NFP (Czaja *et al.*, 2012; Hohnjec *et al.*, 2015). These similarities between the perception and biological activities of NF and Myc-LCOs make it difficult to evaluate the extent to which both sulphated and non-sulphated Myc-LCO root responses may result from inappropriate activation of the NF-signalling pathway in legumes.

In contrast to Myc-LCOs, the Ca²⁺ spiking activity elicited by Myc-COs is unaffected in an *nfp* mutant background (Genre *et al.*, 2013). In addition to the use of whole plants, *M. truncatula* ROCs were also used in these studies since ROCs are readily colonized by AM fungi, but cannot be nodulated and are unresponsive to either rhizobia or NFs. When applied to ROCs Myc-COs are significantly more active in triggering spiking compared to Myc-LCOs (Genre *et al.*, 2013). Together, these results suggest differences in the symbiotic roles played by AM fungal COs and LCOs, and indeed it has been demonstrated that whereas both NF- and Myc-LCOs can stimulate lateral root development (Maillet *et al.*, 2011), this is not the

case for chitotetraose (Olah *et al.*, 2005). Further detailed discussions of these and related findings can be found in the recent reviews of Bucher *et al.*, (2014); Nadal & Paszkowski (2013) and Schmitz & Harrison (2014).

Unfortunately, the absence of AM fungal genetic approaches makes it difficult to ascribe unequivocal signaling roles for either Myc-COs or LCOs. In addition, *nod* gene orthologs were unfortunately not identified in the recently sequenced genome of the AM fungus *Rhizophagus irregularis* (Tisserant *et al.*, 2013). On the other hand, recent studies on non-legume AM hosts are now throwing fresh light on early AM fungal signal perception and in particular the role of certain LysM-RLK receptors. Two research teams have independently shown that CERK1, the rice receptor kinase associated with chitin-triggered immunity is also required for establishing the AM symbiosis. Both knock-out mutant and RNAi experiments have demonstrated that *OsCERK1* is essential for initial AM fungal infection in rice, with a block at the level of epidermal entry (Miyata *et al.*, 2014; Zhang *et al.*, 2015). In contrast, inactivation of OsCEBiP, the second LysM RLK required for perceiving long chain chitin elicitors and activating host immunity, does not result in a defect in AM colonization (Miyata *et al.*, 2014). Interestingly, the closest legume homologs to rice CERK1 are *M. truncatula* LYK3 and *Lotus japonicus* NFR1, both of which are implicated in rhizobial LCO perception. Recent experiments performed using limiting AM inoculation conditions for the two legume hosts have revealed reduced colonization levels for both *Mtlyk3* and *Ljnfr1* mutants (Zhang *et al.*, 2015), thus raising the question of the potential role of these CERK1 orthologs in AM fungal perception in legumes.

In addition to studies on the monocot rice, two additional examples of defective AM phenotypes result from the silencing of LysM-RLKs in non-legume dicots. Firstly, Op den Camp *et al.* (2011) were able to show defects in arbuscule formation in roots of *Parasponia andersonii* following RNAi knockdown of the *PaNFP* gene, although it is not clear whether the initial entry of the AM fungus into the epidermal/cortical tissue is affected. More recently, Buendia *et al.* (2015) have demonstrated by a virus-induced gene silencing approach that knockdown of the *SILYK10* gene, the tomato orthologue of *MtNFP* and *PaNFP*, leads to a block in AM root entry. In conclusion, these important findings in non-legume AM hosts at last provide convincing evidence that fungal symbiotic signals are indeed chitin-based, but at the same time underline the difficulty of deducing the function and precise ligand structure of LysM-RLK receptors based on their phylogenetic proximity.

The fact that non-legume AM hosts such as rice, tomato and carrot do not form additional N-fixing endosymbiotic associations also means that, as for legume ROCs, there should be no interference between AM fungal and rhizobial signaling when studying the activation of the CSSP and associated Ca^{2+} spiking. By introducing a nuclear-localizedameleon into rice, Sun *et al.* (2015) have discovered that Myc-COs (but not Myc-LCOs) are able to elicit Ca^{2+} spiking in rice atrichoblasts, and furthermore at similar concentrations (10^{-8} M) to those used previously for *Medicago* or carrot. Whether the rice *oscerk1* mutant is defective for Myc-CO perception remains to be determined.

III. Pre-infection signaling during *Frankia*/actinorhizal plant nodulation

Gram-positive *Frankia* are filamentous actinomycetes which are able to establish endosymbiotic N-fixing associations with a diverse group of angiosperms (8 plant families and 25 genera) belonging to the Rosid I clade (Santi *et al.*, 2013). These actinorhizal hosts are essentially woody shrubs and trees growing in varied habitats and are natural pioneer species due to their capacity for forming mutually beneficial associations with both *Frankia* and mycorrhizal fungi. Despite major differences between legume and actinorhizal nodule ontogeny and structure, the mechanism of *Frankia* root hair infection is nevertheless highly reminiscent of rhizobial/legume infection. For example, in both *Casuarina* and *Alnus* species, *Frankia* enter the host root *via* infection thread structures formed within root hairs (Wall, 2000). Furthermore, homologs of many components of the CSSP signaling module are present in both actinorhizal hosts (Hocher *et al.*, 2011), and the essential roles of at least two of these (SYMRK and CCaMK) in *Frankia* nodulation has now been clearly demonstrated for *Casuarina glauca* using RNAi approaches (Gherbi *et al.*, 2008; Svistoonoff *et al.*, 2013). On the other hand, little is currently known about the *Frankia* signals that activate this conserved endosymbiotic pathway. Indeed, the absence of the suite of canonical *nod* genes required for NF-like LCO biosynthesis in the sequenced genomes of *Frankia* strains which nodulate *Casuarina* and *Alnus* (Normand *et al.*, 2007), as well as the failure of *Frankia* DNA to complement rhizobial *nod* gene mutants argues that *Frankia*/host recognition involves different types of molecular signals (C  r  monie *et al.*, 1998).

The requirement for a functional CSSP (and in particular CCaMK) in order to initiate *Frankia* infection in actinorhizal hosts also implies that the activation of Ca^{2+} spiking is part of pre-infection *Frankia*-host signaling. This important question has been addressed in two recent publications. Using sonicated extracts from the *Frankia alni* strain ACN14a, Granqvist *et al.*

(2015) were able to show that Ca^{2+} oscillations could be elicited in the root hair cytoplasm of *Alnus glutinosa* after microinjection with calcium dyes. In contrast, Ca^{2+} spiking was not observed in response to NFs of the broad host-range *Rhizobium* NGR234. In a second publication, the expression of the nuclear fluorescent probe (Nup-YC2.1) in transgenic roots of *C. glauca* confirmed that sustained nuclear Ca^{2+} spiking can be triggered following the addition of diluted cell-free *Frankia* supernatants (Chabaud *et al.*, 2015; Fig. 2e). These same supernatants can also activate transcription of the infection-related *CgNIN* gene in *Casuarina* root hairs (Clavijo *et al.*, 2015) and furthermore there is a good correlation between nuclear Ca^{2+} spiking and *promoterCgNIN-GFP* expression as a function of the species-specific *Frankia* strains tested (Chabaud *et al.*, 2015).

Finally, these two pre-infection responses were used as bio-assays for the preliminary characterization of signaling molecules present in the *Frankia* culture supernatants. These experiments revealed that the *Frankia* factors, in contrast to NF LCOs, partition to the aqueous phase after butanol extraction (Chabaud *et al.*, 2015). Furthermore, since chitinase treatment of the *Frankia* supernatant does not abolish either Ca^{2+} spiking or *ProCgNIN* activity, the *Frankia* symbiotic signals are presumably distinct from those of both rhizobia and AM fungi. If so, this also implies that the host receptors which recognize these signals are unlikely to belong to the chitin-binding LysM-RLK family. Thus, a major priority for future research will be to isolate and chemically identify these novel endosymbiotic signaling factors and their corresponding actinorhizal host receptors.

IV. Conclusions & future outlook

Until now, difficulties in manipulating both the actinorhizal woody host plants and the N-fixing actinomycete *Frankia* have retarded research on early pre-infection signaling. However, the development of genetic transformation systems and extensive databases for the *C. glauca* model have at last made it possible to study host responses to secreted *Frankia* factors at the cellular level. Recent results reviewed here now provide final confirmation that nuclear-associated Ca^{2+} oscillatory signaling is indeed a universal hallmark for the activation of the highly conserved CSSP module in endosymbiotic host plants in response to the perception of the appropriate microbial signals. The stage is now set for the purification and chemical characterization of the novel *Frankia* signaling molecules.

Concerning the AM symbiosis, recent successes in identifying infection-defective mycorrhizal phenotypes for certain LysM-RLK mutant or knock-down lines for both rice and

tomato provide a powerful argument in favor of the use of such non-legume AM host species for future studies of both the symbiotic fungal signals and the mechanisms of their perception. In the case of rice, mutants such as *oscerk1*, in combination withameleon-based bioassays for Ca^{2+} signaling/CSSP activation, now offer the means to examine to what extent Myc-CO perception correlates with the AM-defective phenotype, and thus whether short-chain COs may be considered as *bona fide* fungal symbiotic signals.

Acknowledgements

We are grateful to Luisa Lanfranco for critical reading of the manuscript. Financial support was provided to AG by UNITO grant Ricerca Locale 2015 and to MC, DGB by the French National Research Agency grant ANR-12-BSV7-0007-02 and the ‘TULIP’ Laboratory of Excellence grant ANR-10-LABX-41.

References

- Antolin-Llovera M, Ried MK and Parniske M. 2014. Cleavage of the SYMBIOSIS RECEPTOR-LIKE KINASE Ectodomain Promotes Complex Formation with Nod Factor Receptor 5. *Current Biology*, **24**: 422-427.
- Ben Amor B, Shaw SL, Oldroyd GED, Maillet F, Penmetsa RV, Cook D, Long SR, Denarie J and Gough C. 2003. The *NFP* locus of *Medicago truncatula* controls an early step of Nod factor signal transduction upstream of a rapid calcium flux and root hair deformation. *Plant Journal*, **34**: 495-506.
- Bonfante P and Genre A. 2008. Plants and arbuscular mycorrhizal fungi: an evolutionary-developmental perspective. *Trends in Plant Science*, **13**: 492-498.
- Bucher M, Hause B, Krajinski F and Kuester H. 2014. Through the doors of perception to function in arbuscular mycorrhizal symbioses. *New Phytologist*, **204**: 833-840.
- Buendia L, Wang T, Girardin A and Lefebvre B. 2015. The LysM receptor-like kinase SILYK10 regulates the arbuscular mycorrhizal symbiosis in tomato. *New Phytologist*, **210**: 184-195.
- Capoen W, Sun J, Wysham D, Otegui MS, Venkateshwaran M, Hirsch S, Miwa H, Downie JA, Morris RJ, Ane J-M and Oldroyd GED. 2011. Nuclear membranes control symbiotic calcium signaling of legumes. *Proceedings of the National Academy of Sciences of the United States of America*, **108**: 14348-14353.
- C  r  monie H, Debelle F and Fernandez MP. 1999. Structural and functional comparison of *Frankia* root hair deforming factor and rhizobia Nod factor. *Canadian Journal of Botany*, **77**: 1293-1301.
- Chabaud M, Genre A, Sieberer BJ, Faccio A, Fournier J, Novero M, Barker DG and Bonfante P. 2011. Arbuscular mycorrhizal hyphopodia and germinated spore exudates trigger Ca^{2+} spiking in the legume and nonlegume root epidermis. *New Phytologist*, **189**: 347-355.
- Chabaud M, Gherbi H, Pirolles E, Vaissayre V, Fournier J, Moukouanga D, Franche C, Bogusz D, Tisa LS, Barker DG and Svistoonoff S. 2015. Chitinase-resistant hydrophilic symbiotic factors

- secreted by *Frankia* activate both Ca^{2+} spiking and *NIN* gene expression in the actinorhizal plant *Casuarina glauca*. *New Phytologist*, **209**: 86-93.
- Charpentier M and Oldroyd GED. 2013.** Nuclear calcium signaling in plants. *Plant Physiology*, **163**: 496-503.
- Charpentier M, Sun J, Martins TV, Radhakrishnan GV, Findlay K, Soumpourou E, Thouin J, Very A-A, Sanders D, Morris RJ and Oldroyd GED. 2016.** Nuclear-localized cyclic nucleotide-gated channels mediate symbiotic calcium oscillations. *Science*, **352**: 1102-1105.
- Clavijo F, Diedhiou I, Vaissayre V, Brottier L, Acolatse J, Moukouanga D, Crabos A, Auguy F, Franche C, Gherbi H et al. 2015.** The *Casuarina NIN* gene is transcriptionally activated throughout *Frankia* root infection as well as in response to bacterial diffusible signals. *New Phytologist*, **208**: 887-903.
- Czaja LF, Hogeckamp C, Lamm P, Maillet F, Martinez EA, Samain E, Denarie J, Kuester H and Hohnjec N. 2012.** Transcriptional responses toward diffusible signals from symbiotic microbes reveal MtNFP- and MtDMI3-dependent reprogramming of host gene expression by arbuscular mycorrhizal fungal lipochitooligosaccharides. *Plant Physiology*, **159**: 1671-1685.
- Delaux P-M, Radhakrishnan GV, Jayaraman D, Cheem J, Malbreil M, Volkening JD, Sekimoto H, Nishiyama T, Melkonian M, Pokorny L et al. 2015.** Algal ancestors of land plants were preadapted for symbiosis. *Proceedings of the National Academy of Sciences of the United States of America*, **112**: 13390-13395.
- Denarie J and Cullimore J. 1993.** Lipo-oligosaccharide nodulation factors - a new class of signaling molecules mediating recognition and morphogenesis. *Cell*, **74**: 951-954.
- Fournier J, Timmers ACJ, Sieberer BJ, Jauneau A, Chabaud M and Barker DG. 2008.** Mechanism of infection thread elongation in root hairs of *Medicago truncatula* and dynamic interplay with associated rhizobial colonization. *Plant Physiology*, **148**: 1985-1995.
- Genre A, Chabaud M, Balzergue C, Puech-Pages V, Novero M, Rey T, Fournier J, Rochange S, Becard G, Bonfante P and Barker DG. 2013.** Short-chain chitin oligomers from arbuscular mycorrhizal fungi trigger nuclear Ca^{2+} spiking in *Medicago truncatula* roots and their production is enhanced by strigolactone. *New Phytologist*, **198**: 179-189.
- Genre A, Chabaud M, Timmers T, Bonfante P and Barker DG. 2005.** Arbuscular mycorrhizal fungi elicit a novel intracellular apparatus in *Medicago truncatula* root epidermal cells before infection. *Plant Cell*, **17**: 3489-3499.
- Gherbi H, Markmann K, Svistoonoff S, Estevan J, Autran D, Giczey G, Auguy F, Peret B, Laplace L, Franche C et al. 2008.** SymRK defines a common genetic basis for plant root endosymbioses with arbuscular mycorrhiza fungi, rhizobia, and *Frankia* bacteria. *Proceedings of the National Academy of Sciences of the United States of America*, **105**: 4928-4932.
- Granqvist E, Sun J, Op den Camp R, Pujic P, Hill L, Normand P, Morris RJ, Downie JA, Geurts R and Oldroyd GED. 2015.** Bacterial-induced calcium oscillations are common to nitrogen-fixing associations of nodulating legumes and non-legumes. *New Phytologist*, **207**: 551-558.
- Gutjahr C and Parniske M. 2013.** Cell and developmental biology of arbuscular mycorrhiza symbiosis. In *Annual Review of Cell and Developmental Biology*, Vol 29, pp. 593-617.

- 286 **Hocher V, Alloisio N, Auguy F, Fournier P, Doumas P, Pujic P, Gherbi H, Queiroux C, Da Silva**
 287 **C, Wincker P *et al.* 2011.** Transcriptomics of actinorhizal symbioses reveals homologs of the whole
 288 common symbiotic signaling cascade. *Plant Physiology*, **156**: 700-711.
- 289 **Hohnjec N, Czaja-Hasse LF, Hoge Kamp C and Kuester H. 2015.** Pre-announcement of symbiotic
 290 guests: transcriptional reprogramming by mycorrhizal lipochitooligosaccharides shows a strict co-
 291 dependency on the GRAS transcription factors NSP1 and RAM1. *BMC Genomics*, **16**.
- 292 **Kosuta S, Hazledine S, Sun J, Miwa H, Morris RJ, Downie JA and Oldroyd GED. 2008.**
 293 Differential and chaotic calcium signatures in the symbiosis signaling pathway of legumes.
 294 *Proceedings of the National Academy of Sciences of the United States of America*, **105**: 9823-9828.
- 295 **Maillet F, Poinot V, Andre O, Puech-Pages V, Haouy A, Gueunier M, Cromer L, Giraudet D,**
 296 **Formey D, Niebel A *et al.* 2011.** Fungal lipochitooligosaccharide symbiotic signals in arbuscular
 297 mycorrhiza. *Nature*, **469**: 58-63.
- 298 **Miwa H, Sun J, Oldroyd GED and Downie JA. 2006.** Analysis of calcium spiking using aameleon
 299 calcium sensor reveals that nodulation gene expression is regulated by calcium spike number and the
 300 developmental status of the cell. *Plant Journal*, **48**: 883-894.
- 301 **Miyata K, Kozaki T, Kouzai Y, Ozawa K, Ishii K, Asamizu E, Okabe Y, Umehara Y, Miyamoto**
 302 **A, Kobae Y *et al.* 2014.** The bifunctional plant receptor, OsCERK1, regulates both chitin-triggered
 303 immunity and arbuscular mycorrhizal symbiosis in rice. *Plant and Cell Physiology*, **55**: 1864-1872.
- 304 **Nadal M and Paszkowski U. 2013.** Polyphony in the rhizosphere: presymbiotic communication in
 305 arbuscular mycorrhizal symbiosis. *Current Opinion in Plant Biology*, **16**: 473-479.
- 306 **Normand P, Lapierre P, Tisa LS, Gogarten JP, Alloisio N, Bagnarol E, Bassi CA, Berry AM,**
 307 **Bickhart DM, Choise N *et al.* 2007.** Genome characteristics of facultatively symbiotic *Frankia* sp
 308 strains reflect host range and host plant biogeography. *Genome Research*, **17**: 7-15.
- 309 **Olah B, Briere C, Becard G, Denarie J and Gough C. 2005.** Nod factors and a diffusible factor
 310 from arbuscular mycorrhizal fungi stimulate lateral root formation in *Medicago truncatula* via the
 311 DMI1/DMI2 signalling pathway. *Plant Journal*, **44**: 195-207.
- 312 **Oldroyd GED. 2013.** Speak, friend, and enter: signalling systems that promote beneficial symbiotic
 313 associations in plants. *Nature Reviews Microbiology*, **11**: 252-263.
- 314 **Oldroyd GED and Downie JA. 2006.** Nuclear calcium changes at the core of symbiosis signalling.
 315 *Current Opinion in Plant Biology*, **9**: 351-357.
- 316 **Op den Camp R, Streng A, De Mita S, Cao Q, Polone E, Liu W, Ammiraju JSS, Kudrna D,**
 317 **Wing R, Untergasser A *et al.* 2011.** LysM-type mycorrhizal receptor recruited for *Rhizobium*
 318 symbiosis in nonlegume *Parasponia*. *Science (New York, N.Y.)*, **331**: 909-912.
- 319 **Russo G, Spinella S, Sciacca E, Bonfante P and Genre A. 2013.** Automated analysis of calcium
 320 spiking profiles with CaSA software: two case studies from root-microbe symbioses. *BMC Plant*
 321 *Biology*, **13**: 224-236.
- 322 **Santi C, Bogusz D and Franche C. 2013.** Biological nitrogen fixation in non-legume plants. *Annals*
 323 *of Botany*, **111**: 743-767.
- 324 **Schmitz AM and Harrison MJ. 2014.** Signaling events during initiation of arbuscular mycorrhizal
 325 symbiosis. *Journal of Integrative Plant Biology*, **56**: 250-261.

- 326 **Sieberer BJ, Chabaud M, Fournier J, Timmers ACJ and Barker DG. 2012.** A switch in Ca^{2+}
 327 spiking signature is concomitant with endosymbiotic microbe entry into cortical root cells of
 328 *Medicago truncatula*. *Plant Journal*, **69**: 822-830.
- 329 **Sieberer BJ, Chabaud M, Timmers AC, Monin A, Fournier J and Barker DG. 2009.** A nuclear-
 330 targetedameleon demonstrates intranuclear Ca^{2+} spiking in *Medicago truncatula* root hairs in
 331 response to rhizobial nodulation factors. *Plant Physiology*, **151**: 1197-1206.
- 332 **Singh S, Katzer K, Lambert J, Cerri M and Parniske M. 2014.** CYCLOPS, A DNA-binding
 333 transcriptional activator, orchestrates symbiotic root nodule development. *Cell Host & Microbe*, **15**:
 334 139-152.
- 335 **Sun J, Miller JB, Granqvist E, Wiley-Kalil A, Gobbato E, Maillet F, Cottaz S, Samain E,**
 336 **Venkateshwaran M, Fort S et al. 2015.** Activation of symbiosis signaling by arbuscular mycorrhizal
 337 fungi in legumes and rice. *Plant Cell*, **27**: 823-838.
- 338 **Svistoonoff S, Benabdoun FM, Nambiar-Veetil M, Imanishi L, Vaissayre V, Cesari S, Diagne N,**
 339 **Hocher V, de Billy F, Bonneau J et al. 2013.** The independent acquisition of plant root nitrogen-
 340 fixing symbiosis in fabids recruited the same genetic pathway for nodule organogenesis. *Plos One*, **8**:
 341 e64515.
- 342 **Tisserant E, Malbreil M, Kuo A, Kohler A, Symeonidi A, Balestrini R, Charron P, Duensing N,**
 343 **Frey NFD, Gianinazzi-Pearson V et al. 2013.** Genome of an arbuscular mycorrhizal fungus provides
 344 insight into the oldest plant symbiosis. *Proceedings of the National Academy of Sciences of the United*
 345 *States of America*, **110**: 20117-20122.
- 346 **Venkateshwaran M, Jayaraman D, Chabaud M, Genre A, Balloon AJ, Maeda J, Forshey K, den**
 347 **Os D, Kwiecien NW, Coon JJ et al. 2015.** A role for the mevalonate pathway in early plant symbiotic
 348 signaling. *Proceedings of the National Academy of Sciences of the United States of America*, **112**:
 349 9781-9786.
- 350 **Venkateshwaran M, Volkening JD, Sussman MR and Ane J-M. 2013.** Symbiosis and the social
 351 network of higher plants. *Current Opinion in Plant Biology*, **16**: 118-127.
- 352 **Wall LG. 2000.** The actinorhizal symbiosis. *Journal of Plant Growth Regulation*, **19**: 167-182.
- 353 **Zhang X, Dong W, Sun J, Feng F, Deng Y, He Z, Oldroyd GED and Wang E. 2015.** The receptor
 354 kinase CERK1 has dual functions in symbiosis and immunity signalling. *Plant Journal*, **81**: 258-267.

Figure Legends

Figure 1

Schema illustrating the role of the Common Symbiosis Signaling Pathway (CSSP)

A number of plant genes and secondary messengers are required for the successful functioning of the conserved CSSP core module, first discovered in the model legume species *Lotus japonicus* and *Medicago truncatula*, and now extended to plant hosts of all known root endosymbiotic associations including the rhizobial/legume, AM and actinorhizal symbioses. Although several important features of this signaling module still remain to be elucidated, we present here a consensus linear representation of the best-studied pathway components which have been identified for both model legumes, organized around the activation of the nuclear-associated Ca^{2+} spiking response observed in cells of the root epidermis (root hairs or atrichoblasts) prior to the initiation of apoplastic microbial infection (see text). In the case of the rhizobial and AM symbioses the CSSP is activated following symbiotic signal perception by plasma membrane (PM) localized LysM-RLK receptors, most probably part of a larger complex including the leucine-rich repeat receptor-like kinase known as LjSYMRK/MtDMI2 (Antolin-Llovera *et al.*, 2014). SYMRK can also interact with 3-hydroxy-3-methylglutaryl-CoA reductase (HMGR), a key enzyme in the so-called mevalonate pathway, a source of potential secondary messengers including mevalonate itself (Venkateshwaran *et al.*, 2015). Following signal transduction from the PM to the nucleus, nuclear membrane cation channels known as LjCASTOR/LjPOLLUX/MtDMI1, likely in association with the recently discovered cyclic nucleotide gated-calcium channel complex CNGC15 (Charpentier *et al.*, 2016) are then required for rapid Ca^{2+} release and the initiation of nucleoplasmic Ca^{2+} spiking. Efficient re-uptake of Ca^{2+} across the nuclear membrane between repeated spiking requires a calcium ATPase pump which has been identified as MCA8 in *M. truncatula* (Capoen *et al.*, 2011). The subsequent decoding of the intranuclear Ca^{2+} oscillatory response involves two key associated components (LjCCaMK/MtDMI3 and LjCYCLOPS/MtIPD3). Binding of Ca^{2+} to the calcium and calmodulin-dependent kinase CCaMK (both directly and indirectly via Ca^{2+} /calmodulin) leads to phosphorylation of the coiled-coil protein CYCLOPS (Singh *et al.*, 2014). Finally, the activation of a downstream signaling cascade *via* a repertoire of GRAS/ERF transcription factors results in the synthesis of the suite of proteins required for the transcriptional re-modeling of the epidermal cell in preparation for apoplastic infection. Note that only recent references have been included here, and that further details about the CSSP and nuclear-associated Ca^{2+} spiking can be found in a number of comprehensive review

articles (e.g. Charpentier & Oldroyd, 2013; Gutjahr & Parniske, 2013; Oldroyd, 2013; Venkateshwaran *et al.*, 2013). Note also that in the case of the N-fixing actinorhizal association, the nature of both the *Frankia* signal (AF=Actinorhizal Factor) and the corresponding host receptor remain to be determined, and that direct evidence for an essential role in microbial/host signaling has only been demonstrated so far for *CgSYMRK* and *CgCCaMK* (see text).

Figure 2

Nuclear-associated Ca²⁺ spiking in response to bacterial and fungal symbiotic signals

(a-c). Nuclear Ca²⁺ responses to AM fungal symbiotic factors were recorded in epidermal atrichoblasts using *Medicago truncatula* (Mt) root organ cultures (ROCs) expressing the Nup-YC2.1ameleon (Chabaud *et al.*, 2011; Genre *et al.*, 2013). Representative spiking observed over a 20 min period are shown for (a) cells in direct contact with a fungal hyphopodium prior to infection, or following treatment with either (b) a germinated spore exudate (concentrated 10-fold) of *Gigaspora rosea* or (c) a 10⁻⁸ M solution of chitotetraose (CO4). (d,e). The identicalameleon reporter was used to compare the nuclear Ca²⁺ spiking observed in (d) *M. truncatula* root hairs (intact plant) treated with 10⁻⁹ M *Sinorhizobium meliloti* (Sm) Nod factor with (e) *Casuarina glauca* (Cg) root hair spiking in response to a crude *Frankia* Cci3 supernatant (SN, diluted 100-fold) (Chabaud *et al.*, 2015). Note that the spiking patterns in response to AM fungal signaling in atrichoblasts are less regular in both frequency and individual spike profile as compared to either *Rhizobium* or *Frankia*-elicited spiking in host root hairs.




Figure 1




Figure 2